

**WISCONSIN**

Information Retrieval  
For Business and Industry

**TECH SEARCH**

WTS Number: 444575



Request Date: 2/14/07 1:27 PM

Conf Number: 141466

Requester: Lee R. Nemchek

Morrison & Foerster LLP  
555 W. Fifth Street  
Suite 3500  
Los Angeles, CA 90013

Need by:

02/16/2007 - Standard

Company Phone:

Requester Phone: 213-892-5359

Fax:

Requester Email: lnemchek@mofo.com

Send-To Email: lnemchek@mofo.com

Reference: 51471-20029.00-09185

Delivery: Email

Instructions:

11. Grohmann, K. et al. (2001). "Mutations in the Gene Encoding Immunoglobulin mu-Binding Protein 2 Cause Spinal Muscular Atrophy with Respiratory Distress Type 1," Nat. Genet. 29:75-77.

*Prim*  
*Med*  
*Q7*  
*N2563*

An outreach service of the Kurt F. Wendt Library, University of Wisconsin - Madison  
Email: wts@engr.wisc.edu | Web: <http://www.wisc.edu/techsearch> | Phone: (608) 262-5917

Copyright Royalty: \$ \_\_\_\_\_

Refer Off Campus

# Of Pages: \_\_\_\_\_

1061-4K36

# nature *genetics*

ASL

volume 29 no. 1

september 2001

HEALTH SCIENCES LIBRARY  
UNIVERSITY OF WISCONSIN

SEP 17 2001

1905 Linden Drive  
Madison, WI 53706

Policing p53

Archaeal legacy

Insulin insights

Translating  
spermatogenesis



<http://genetics.nature.com>



Cover art by:  
Michael Malicki  
(tempera on paper)

# nature genetics

volume 29 no. 1

september 2001

## editorial

- Something old, something new 1

## news & views

- A new verdict for an old convict 3  
Gerd P Pfeifer • SEE ALSO 25
- The power of public access 4  
John Quackenbush • SEE ALSO 88
- Choreographing mRNA biogenesis 6  
Charles N Cole
- The sights along route 65 8  
John C Saari • SEE ALSO 70
- TOUCHINGbase 11

## book review

- Genetics in primary care: what do we expect? 13  
Charles J Epstein

## correspondence

- How human geneticists in US view commercialization of  
the Human Genome Project 15  
I Rabino

## brief communications

- Mutations in *SEPN1* cause congenital muscular dystrophy  
with spinal rigidity and restrictive respiratory syndrome 17  
B Moghadaszadeh, N Petit, C Jaillard, M Brockington, S Q Roy, L Merlini, N Romero,  
B Estournet, I Desguerre, D Chaigne, F Muntoni, H Topaloglu & P Guicheney

Nature Genetics (ISSN 1061-4036) is published monthly by Nature America Inc., headquartered at 345 Park Avenue South, New York, NY 10010-1707. Editorial Office: 345 Park Avenue South, New York, NY 10010-1707. Tel: (212) 726 9314, Fax: (212) 545 8341. North American Advertising: 345 Park Avenue South, New York, NY 10010-1707. Tel: (212) 726 9200, Fax: (212) 696 9006. European Advertising: Nature Genetics, Porters South, Crinan Street, London N1 9XW, UK. Tel: 44 171 833 4000, Fax: 44 171 843 4996. New subscriptions, renewals, changes of address, back issues and all customer service questions in the Americas should be addressed to: Nature Genetics Subscription Department, P.O. Box 5054, Brentwood, TN 37024-5054. Tel: (800) 524 0384, Direct Dial (615) 377 3322, Fax: (615) 377 0525. Outside the Americas: Nature Genetics, Macmillan Magazines Ltd., Houndsmill, Brunel Road, Basingstoke, RG21 6XS, U.K. Tel: +44-(0)1256-329242. Fax: +44-(0)1256 812358. Email: subscriptions@nature.com. Annual subscription rates: North America: US\$650 (institutional/corporate), US\$199 (individual making personal payment), Canada add 7% for GST, BN: 14091 1595 RT; U.K./Europe: £475 (institutional/corporate), £185 (individual making personal payment), £99 (student); Rest of world (excluding Japan): £520 (institutional/corporate), £235 (individual making personal payment) £110 (student); Japan: Contact Nature Japan K.K., Shin-Mitsuke Building 3F, 3-6 Ichigaya Tamachi, Shinjuku-ku, Tokyo 162-0843. Periodicals postage paid at New York, NY 10010, and at additional mailing offices. POSTMASTER: Send address changes to Nature Genetics Subscription Department, P.O. Box 5054, Brentwood, TN 37024-5054. Executive Officers of Nature America Inc.: Annette Thomas, President; Edward Valis, Secretary & Treasurer. Published in Japan by Nature Japan K.K., Shin-Mitsuke Building 3F, 3-6 Ichigaya Tamachi, Shinjuku-ku, Tokyo 162-0843, Japan. Printed by Cadmus Journal Services, Richmond, VA, US. Copyright ©2001 Nature America Inc.

# Mutations in the gene encoding immunoglobulin $\mu$ -binding protein 2 cause spinal muscular atrophy with respiratory distress type 1

Katja Grohmann<sup>1</sup>, Markus Schuelke<sup>1</sup>, Alexander Diers<sup>1</sup>, Katrin Hoffmann<sup>2</sup>, Barbara Lucke<sup>2</sup>, Coleen Adams<sup>3</sup>, Enrico Bertini<sup>4</sup>, Hajnalka Leonhardt-Horti<sup>5</sup>, Francesco Muntoni<sup>6</sup>, Robert Ouvrier<sup>7</sup>, Arne Pfeufer<sup>8</sup>, Rainer Rossi<sup>9</sup>, Lionel Van Maldergem<sup>10</sup>, Jo M. Wilmschurst<sup>7</sup>, Thomas F. Wienker<sup>11</sup>, Michael Sendtner<sup>12</sup>, Sabine Rudnik-Schöneborn<sup>13</sup>, Klaus Zerres<sup>13</sup> & Christoph Hübner<sup>1</sup>

Published online: 13 August 2001, DOI: 10.1038/ng703

Classic spinal muscular atrophy (SMA) is caused by mutations in the telomeric copy of *SMN1*. Its product is involved in various cellular processes, including cytoplasmic assembly of spliceosomal small nuclear ribonucleoproteins, pre-mRNA processing and activation of transcription<sup>1–3</sup>. Spinal muscular atrophy with respiratory distress (SMARD) is clinically and genetically distinct from SMA<sup>9–13</sup>. Here we demonstrate that SMARD type 1 (SMARD1) results from mutations in the gene encoding immunoglobulin  $\mu$ -binding protein 2 (*IGHMBP2*; on chromosome 11q13.2–q13.4). In six SMARD1 families, we detected three recessive missense mutations (exons 5, 11 and 12), two nonsense mutations (exons 2 and 5), one frameshift deletion (exon 5) and one splice donor-site mutation (intron 13). Mutations in mouse *Ighmbp2* (ref. 14) have been shown to be responsible for spinal muscular atrophy in the neuromuscular degeneration (*nmd*) mouse<sup>15</sup>, whose phenotype resembles the SMARD1 phenotype. Like the *SMN1* product, *IGHMBP2* colocalizes with the RNA-processing machinery in both the cytoplasm and the nucleus<sup>16–19</sup>. Our results show that *IGHMBP2* is the second gene found to be defective in spinal muscular atrophy, and indicate that *IGHMBP2* and *SMN* share common functions important for motor neuron maintenance and integrity in mammals.

Autosomal recessive SMARD (also known as diaphragmatic spinal muscular atrophy<sup>11,13</sup>, distal hereditary motor neuropathy type VI, dHMN-VI (ref. 20) and severe infantile axonal neuropathy with respiratory failure<sup>21</sup>) and classic autosomal recessive SMA are both characterized by dysfunction and progressive loss of  $\alpha$ -motor neurons in the anterior horn of the spinal cord, leading to neurogenic muscular atrophy with subsequent symmetrical muscle weakness of trunk and limbs<sup>9–13</sup>. In contrast to SMA, distal muscles are more severely affected in SMARD, and life-threatening respiratory distress with clinical and radiological evidence of unilateral or bilateral paralysis of the diaphragm is the most prominent presenting symptom<sup>9–13</sup> (Fig. 1 and Table 1). In previous studies, clinical and genetic heterogeneity of SMARD has been demonstrated. SMARD type 1 with non-congenital onset of respiratory distress has been linked to chromosome 11q13–q21 (SMARD1), whereas linkage to this locus could be excluded in one family with two affected children suffering from respiratory distress of congenital onset<sup>12</sup>.

Symptoms similar to those of human SMARD1 have been found in *nmd* mice homozygous for autosomal recessive *Ighmbp2* mutations<sup>14</sup>. These animals suffer from progressive paralysis of the limbs with onset at 2 weeks of age, leading to death by 3.5 weeks secondary to respiratory failure<sup>14,15</sup>. Histopathological analysis showed progressive degeneration of  $\alpha$ -motor neurons with secondary generalized atrophy of distal limb muscles<sup>15</sup>.

We have refined the SMARD1 locus to a genetic interval of 9 cM with the centromeric critical breakpoint distal to *D11S913* and the telomeric breakpoint proximal to *D11S916* (data not shown). The most promising candidate gene within this region was *IGHMBP2*, as mutations of the mouse ortholog are responsible for the *nmd* phenotype<sup>14</sup>. Human *IGHMBP2* is composed of 15 exons (for exon–intron boundaries, see Web Table A). It is ubiquitously expressed<sup>22</sup>, with the highest levels of *IGHMBP2* mRNA detected in testis and low-to-moderate expression in other human tissues<sup>23</sup>.



Fig. 1 Chest X-ray showing eventration of the right hemidiaphragm in a SMARD1 patient at 8 weeks of age. The infant presented with severe respiratory distress resulting from paralysis of the diaphragm.

<sup>1</sup>Department of Neuropediatrics, Charité, Campus Virchow-Klinikum, Humboldt University, 13353 Berlin, Germany. <sup>2</sup>Gene Mapping Center, Max Delbrueck Center for Molecular Medicine, 13092 Berlin-Buch, Germany. <sup>3</sup>Division of Pediatric Neurology, Alberta Children's Hospital, University of Calgary, Calgary, Alberta, T2T 5C7, Canada. <sup>4</sup>Department of Neurosciences and Unit of Molecular Medicine, Bambino Gesù Children's Hospital, 00165 Rome, Italy. <sup>5</sup>Children's Hospital, 63069 Offenbach, Germany. <sup>6</sup>Department of Paediatrics, Hammersmith Hospital, London W12 0HS, UK. <sup>7</sup>Institute for Neuromuscular Research, Children's Hospital at Westmead, Parramatta, NSW 2124, Australia. <sup>8</sup>Institute of Human Genetics, GSF Research Institute, 85764 Neuherberg, Germany. <sup>9</sup>Children's Hospital Neukölln, 12051 Berlin, Germany. <sup>10</sup>Centre de Génétique Humaine, Institut de Pathologie et de Génétique, 6280 Lovreval, Belgium. <sup>11</sup>Institute of Medical Biometry, University of Bonn, 53105 Bonn, Germany. <sup>12</sup>Institute of Clinical Neurobiology, University of Würzburg, 97080 Würzburg, Germany. <sup>13</sup>Department of Human Genetics, Technical University, 52074 Aachen, Germany. Correspondence should be addressed to C.H. (e-mail: christoph.huebner@charite.de).

Table 1 • Mutations of *IGHMBP2* and clinical data in SMARD1 patients

Family	Affected/unaffected sibs	Geographic origin	Consanguinity	Homozygous (heterozygous) mutation	Location	Amino acid substitution	Class of mutation	Age at onset of respiratory distress (weeks)
1	1/1	South Italy	no	1540G→A	exon 11	E514K	missense	4
2	5/3	Lebanon	yes	638A→G	exon 5	H213R	missense	6-9 (family 1 in ref. 12)
3	1/2	Turkey	yes	1738G→A	exon 12	V580I	missense	8
4	2/1	Germany	no	(121C→T) (675delT)	exon 2 exon 5	Q41X -	nonsense frameshift	9-12 (family 2 in ref. 12)
5	1/1	Lebanon	yes	707T→G	exon 5	L236X	nonsense	16 (patient 3 in ref. 21)
6	1/1	Sicily	yes	IVS13+1G→T	intron 13	-	splice donor	18

Family numbers correspond to those in Fig. 3.

DNA sequence analysis of four consanguineous families (families 2, 3, 5 and 6) and two non-consanguineous families (families 1 and 4) demonstrated seven different *IGHMBP2* mutations in four different exons and in one intron (Table 1). In family 1, a homozygous 1540G→A transition in exon 11 predicts substitution of a glutamic acid by lysine (E514K), and in family 2, a homozygous 638A→G transition in exon 5 predicts substitution of a histidine by arginine (H213R). This histidine residue is located within the first of seven helicase domains<sup>24</sup>. A homozygous 1738G→A transition in exon 12 (family 3) predicts replacement of valine by isoleucine (V580I). The residues E514, H213 and V580 affected by the three missense mutations are conserved between the orthologs of man, mouse, rat and golden hamster (Fig. 2; Web Fig. A). In addition, we detected a heterozygous nonsense mutation in exon 2 (Q41X; family 4) and a homozygous nonsense mutation in exon 5 (L236X; family 5). In family 4, the second allele carried a 1-bp deletion in exon 5 (675delT), resulting in a nonsense peptide of six amino acids after V225, with subsequent chain termination. In family 6, a homozygous point mutation at the consensus splice donor site of intron 13 (IVS13+1G→T) probably results in defective splicing. Neither the splice donor-site mutation nor the missense mutations were detected in 50 unaffected unrelated individuals, indicating that these mutations do not reflect common polymorphisms. The *IGHMBP2* mutations segregate with the disease phenotype in all

families (Fig. 3). This is consistent with the proposed autosomal recessive mode of inheritance<sup>12</sup> and supports the hypothesis that the mutations cause SMARD1.

We found a silent 180C→T sequence variation in exon 2 (Y60Y; family 2) and detected several variations in both affected people and healthy controls. These variations would seem to be polymorphisms (5' untranslated region-136insGCCTCTTCCCCG, families 3 and 6; 2636C→A, T879K, families 2, 4 and 5; IVS14+54G→A, families 2 and 5). *IGHMBP2* consists of 993 amino acids and includes 7 putative helicase motifs<sup>24</sup> and a DEAD box-like motif, which is typical for RNA helicases<sup>17</sup>. *IGHMBP2* contains a DNA-binding domain at position 638-786 including the helicase motifs V and VI (refs. 19,22,24) and the nucleic acid-binding R3H motif<sup>25</sup>. The cellular function of *IGHMBP2* is unknown. It is involved in




Fig. 2 Alignment of selected regions of human *IGHMBP2* with orthologs of other species. Arrows indicate positions of the missense and nonsense mutations in SMARD1 patients. Family numbers correspond to those in Table 1. f1-5, families 1-5. (For alignment of the whole amino-acid sequences, see Web Fig. A).




Fig. 3 Segregation of *IGHMBP2* mutations (restriction fragment-length polymorphism analysis). In family 4 (f4), the two affected siblings were compound heterozygotes carrying the maternal 121C→T nonsense mutation and the paternal 675delT frameshift deletion. Family numbers correspond to those in Table 1. co, control; f1-6, families 1-6; mt, mutated *IGHMBP2*; wt, wildtype *IGHMBP2*.

immunoglobulin-class switching<sup>22</sup>, in pre-mRNA processing<sup>17</sup> and in regulation of transcription by DNA binding<sup>16,19</sup> or by interacting with TATA-binding protein<sup>18</sup>. In this respect, IGHMBP2 resembles the SMN protein, which is defective in classic SMA. SMN binds directly to DNA and RNA (ref. 26), activates transcription by association with the viral nuclear transcription activator E2 (ref. 4) and is involved in pre-mRNA processing<sup>6,8</sup>. Although SMN does not contain helicase motifs, it is part of a major cellular complex including DP103, a member of the DEAD box family of RNA helicases<sup>3,5,27</sup>.

Our findings support the hypothesis that mutant SMN and mutant IGHMBP2 result in a similar dysfunction of spinal motor neurons, resulting in SMA and SMARD1, respectively. Functional characterization of IGHMBP2 will help to unravel the enigma of the cellular processes that underlie the specificity of diseases leading to neurogenic muscular atrophy.

## Methods

**Patients.** We studied a total of 11 patients and 21 relatives from 6 unrelated families (Table 1). The diagnosis of SMARD1 was made on the basis of clinical criteria<sup>11-13</sup>. All affected infants were floppy, presented with life-threatening respiratory distress and had unilateral or bilateral eventration of the diaphragm on chest X-ray (Fig. 1). Surviving patients were on long-term artificial ventilation. In addition, analysis of muscle biopsy specimen showed neurogenic muscular atrophy. One patient (family 5) had bilateral equinovarus foot deformities at birth<sup>21</sup>. In all families, haplotype analysis was consistent with linkage to 11q13. We obtained blood samples from patients and family members after obtaining informed consent according to the declaration of Helsinki. We isolated DNA from peripheral blood lymphocytes, Guthrie card samples and skin fibroblast cultures according to standard procedures.

**Haplotype analysis.** We used 12 fluorescently labeled polymorphic markers and standard semi-automated methods<sup>12</sup> for microsatellite analysis. We used a MegaBACE 1000 DNA-sequencer and markers from the Génethon final linkage map: D11S1883, D11S1913, D11S4095, D11S4178, D11S1314, D11S1916, D11S1901, D11S1358, D11S1311, D11S4176, D11S1757 and D11S1917.

**Database analysis.** The mRNA sequence of IGHMBP2 has been published, and we used homology search and exon assembly with BLAST programs at the National Center for Biotechnology Information. We derived the genomic sequence of IGHMBP2 from a contig of three genomic clones.

**Sequence analysis.** We amplified all 15 exons of IGHMBP2 from patients' genomic DNA with intronic primers (see Web Table B). We used standard procedures for bi-directional automatic sequencing with fluorescent dye terminators on the MegaBACE 1000 DNA-sequencer with the above-mentioned primers.

We verified the intrafamilial segregation of the mutations by restriction fragment-length polymorphism analysis (Fig. 3). When we found no natural restriction sites, we used primer-induced restriction analysis (primer mismatches underlined): 1540G→A, 11F/5'-CCTGGATGTGCAAACTGACGA.GGCGGACGT-3' (AatII, mutated (mt)=124 bp, wildtype (wt)=98+26 bp); 638A→G, 5F/5R (NcoI, mt=255 bp, wt=149+105 bp); 1738G→A, 12F/5'-GGCTCCGTACCTTCTCTGTGGATCTCA-3' (BspHI, mt=211+30 bp, wt=241 bp); 121C→T, 2F/2R (XbaI, mt=92+199 bp, wt=291 bp); 675delT, 5F/5'-CCTTGTTCACAGCTTGAAGAATGATGTCA-3' (HincII, mt=216 bp, wt=188+29 bp); 707T→G, 5F/5'-TGGCATGCACTGCCAC.CCTT-3' (AflIII, mt=241 bp, wt=211+30 bp); IVS13+1G→T, 13.2F/5'-CACTGCCCAAGTCTTATTAGTTGAGTTA-3' (HpaI, mt=277+29 bp, wt=306 bp).

**Accession numbers.** GenBank: IGHMBP2, L14754; IGHMBP2 genomic clones, AP000808.2, AP000444.3, AC019166.5; *Ighmbp2* rat (*Rattus Norvegicus*), AF199411; OMIM: SMA1, #253300; SMARD1, #604320. SwissProt: IGHMBP2 human (*Homo Sapiens*), P38935; *Ighmbp2* mouse (*Mus Musculus*), P40694; *Ighmbp2* hamster (*Meocricetus Auratus*), Q60560.

**Note:** Supplementary information is available on the Nature Genetics web site ([http://genetics.nature.com/supplementary\\_info/](http://genetics.nature.com/supplementary_info/)).

## Acknowledgments

We thank the patients and their families for participation in the study. The help, discussions and critical comments on this manuscript of C. Bassir, M. Bollinger, S. Buttner, E. Eike, A. Huebner, A.Y. Manzur, J. Scholz, G. Stoltenburg-Didinger and A. Zwirner are acknowledged. This study has been supported by grants from the Deutsche Forschungsgemeinschaft (Hu 408/3-1, K.G. and C.H.; Ze 205/10-1, S.R.-S. and K.Z.; SFB 581, TPB1, M.S., Würzburg) and in part by the parents' support group 'Helft dem muskelkranken Kind', Hamburg, Germany (C.H.).

Received 14 June; accepted 5 July 2001.

- Fischer, U., Liu, Q. & Dreyfuss, G. The SMN-SIP1 complex has an essential role in spliceosomal snRNP biogenesis. *Cell* 90, 1023-1029 (1997).
- Lefebvre, S., Bürglen, L., Frézal, J., Munnich, A. & Melki, J. The role of the SMN gene in proximal spinal muscular atrophy. *Hum. Mol. Genet.* 7, 1531-1536 (1998).
- Charroux, B. et al. Gemin3: a novel DEAD box protein that interacts with SMN, the spinal muscular atrophy gene product, and is a component of gems. *J. Cell Biol.* 147, 1181-1193 (1999).
- Strasswimmer, J. et al. Identification of survival motor neuron as a transcriptional activator-binding protein. *Hum. Mol. Genet.* 8, 1219-1226 (1999).
- Campbell, L. et al. Direct interaction of Smn with dp103, a putative RNA helicase: a role for Smn in transcription regulation? *Hum. Mol. Genet.* 9, 1093-1100 (2000).
- Meister, G. et al. Characterization of a nuclear 20S complex containing the survival of motor neurons (SMN) protein and a specific subset of spliceosomal Sm proteins. *Hum. Mol. Genet.* 9, 1977-1986 (2000).
- Jablonska, S. et al. Co-regulation of survival motor neuron (SMN) protein and its interactor SIP1 during development and in spinal muscular atrophy. *Hum. Mol. Genet.* 10, 497-505 (2001).
- Pellizzoni, L., Charroux, B., Rappsilber, J., Mann, M. & Dreyfuss, G. A functional interaction between the survival motor neuron complex and RNA polymerase II. *J. Cell Biol.* 152, 75-85 (2001).
- Mellins, R.B., Hays, A.P., Gold, A.P., Berdon, W.E. & Bowdler, J.D. Respiratory distress as the initial manifestation of Werdnig-Hoffmann disease. *Pediatrics* 53, 33-40 (1974).
- Bertini, E. et al. Distal infantile spinal muscular atrophy associated with paralysis of the diaphragm: a variant of infantile spinal muscular atrophy. *Am. J. Med. Genet.* 33, 328-335 (1989).
- Rudnik-Schöneborn, S., Forkert, R., Hahnen, E., Wirth, B. & Zerres, K. Clinical spectrum and diagnostic criteria of infantile spinal muscular atrophy: further delineation on the basis of SMN gene deletion findings. *Neuropediatrics* 27, 8-15 (1996).
- Grohmann, K. et al. Diaphragmatic spinal muscular atrophy with respiratory distress is heterogeneous, and one form is linked to chromosome 11q13-q21. *Am. J. Hum. Genet.* 65, 1459-1462 (1999).
- Zerres, K. & Davies, K.E. 59th ENMC International Workshop: Spinal Muscular Atrophies: recent progress and revised diagnostic criteria 17-19 April 1998, Soestduinen, The Netherlands. *Neuromuscul. Disord.* 9, 272-278 (1999).
- Cox, G.A., Mahaffey, C.L. & Frankel, W.N. Identification of the mouse neuromuscular degeneration gene and mapping of a second site suppressor allele. *Neuron* 21, 1327-1337 (1998).
- Cook, S.A., Johnson, K.R., Bronson, R.T. & Davisson, M.T. Neuromuscular degeneration (*nmd*): a mutation on mouse chromosome 19 that causes motor neuron degeneration. *Mamm. Genome* 6, 187-191 (1995).
- Chen, N.N., Kerr, D., Chang, C.-F., Honjo, T. & Khalili, K. Evidence for regulation of transcription and replication of the human neurotropic virus JCV genome by the human Subp-2 protein in glial cells. *Gene* 185, 55-62 (1997).
- Molnar, G.M. et al. Association of the mammalian helicase MAH with the pre-mRNA splicing complex. *Proc. Natl. Acad. Sci. USA* 94, 7831-7836 (1997).
- Zhang, Q., Wang, Y.-C.J. & Mohtai, E.A. Subp-2 represses the Epstein-Barr virus lytic switch promoter. *Virology* 255, 160-170 (1999).
- Miao, M., Chan, S.-L., Fletcher, G.L. & Hew, C.L. The rat ortholog of the presumptive flounder antifreeze enhancer-binding protein is a helicase domain-containing protein. *Eur. J. Biochem.* 267, 7237-7245 (2000).
- McEntagart, M. et al. Localization of the gene for distal hereditary motor neuropathy VII (dHMN-VII) to chromosome 2q14. *Am. J. Hum. Genet.* 68, 1270-1276 (2001).
- Wilmshurst, J.M. et al. Severe infantile axonal neuropathy with respiratory failure. *Muscle Nerve* 24, 760-768 (2001).
- Fukita, Y. et al. The human Subp-2, a DNA-binding protein specific to the single-stranded guanine-rich sequence related to the immunoglobulin  $\mu$  chain switch region. *J. Biol. Chem.* 268, 17463-17470 (1993).
- Mohan, W.S. et al. Human S mu binding protein-2 binds to the drug response element and transactivates the human apoA-I promoter: role of gemfibrozil. *J. Lipid Res.* 39, 255-267 (1998).
- Mizuta, T.-R., Fukita, Y., Miyoshi, T., Shimizu, A. & Honjo, T. Isolation of cDNA encoding a binding protein specific to 5'-phosphorylated single-stranded DNA with G-rich sequences. *Nucleic Acids Res.* 21, 1761-1766 (1993).
- Grishin, N.V. The R3H motif: a domain that binds single-stranded nucleic acids. *Trends Biochem. Sci.* 23, 329-330 (1998).
- Lorson, C.L. & Androphy, E.J. The domain encoded by exon 2 of the survival motor neuron protein mediates nucleic acid binding. *Hum. Mol. Genet.* 7, 1269-1275 (1998).
- Grundhoff, A.T. et al. Characterization of DP103, a novel DEAD box protein that binds to the Epstein-Barr virus nuclear proteins EBNA2 and EBNA3C. *J. Biol. Chem.* 274, 19136-19144 (1999).